Zinner syndrome: an updated pooled analysis based on 214 cases from 1999 to 2020: systematic review

Tianzhu Liu^{1#}, Xiaodan Li^{2#}, Lesheng Huang¹, Hongyi Li¹, Kaili Cai¹, Jinghua Jiang¹, Nianli Chen³, Wanchun Zhang¹, Jiahui Tang¹, Meng Zhang¹, Dan Zhao¹, Jingzhi Ye¹, Chengfeng Zhang¹, Tao He¹, Bo Liu¹, Yongsong Ye¹, Jun Chen¹

¹Department of Radiology, Guangdong Hospital of Traditional Chinese Medicine, Zhuhai, China; ²Department of Gynecology, Guangdong Hospital of Traditional Chinese Medicine, Zhuhai, China; ³Department of Urology, Guangdong Hospital of Traditional Chinese Medicine, Zhuhai, China

Contributions: (I) Conception and design: B Liu, Y Ye, J Chen; (II) Administrative support: J Tang, J Ye, C Zhang, T He; (III) Provision of study materials or patients: H Li, L Huang; (IV) Collection and assembly of data: K Cai, J Jiang, N Chen, W Zhang; (V) Data analysis and interpretation: T Liu, X Li, M Zhang, D Zhao; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work.

Correspondence to: Jun Chen; Bo Liu; Yongsong Ye. Department of Radiology, Guangdong Hospital of Traditional Chinese Medicine, No. 111 Dade Road, Zhuhai 519000, China. Email: junesums@163.com; liubogzcm@163.com; ye7yong7song@163.com.

Abstract: Zinner syndrome (ZS) is a rare anomaly of the Wolffian duct. We searched for case reports and case series to identify the most effective methods for examination and treatment. The PubMed/Medline, Embase, and Scopus databases were searched. Our searches yielded 160 case reports/case series and included 214 patients. The mean age at diagnosis was 29.35 years. The R:L distribution was 1:1. The most common symptoms were: frequency (24.3%), dysuria (23.1%) and perineal pain (20.2%) in ZS patients with clinical symptoms. The diagnostic investigations used most frequently were magnetic resonance imaging (MRI) (67.8%) and ultrasonography (65.0%). Maximum flow rate on uroflowmetry, semen ejaculate volume, sperm count, and sperm motility were significantly lower in patients with ZS. Among 193 patients with treatment details, 65.8% underwent surgery; 9.8% underwent aspiration; 24.3% were followed with observation. Two patients died after surgery; all other patients remained asymptomatic after surgery. Fifty-two patients had complications and comorbidities. In patients with ZS, the most common symptoms are seminal vesicle cyst enlargement and compression of the bladder, ureter, and reproductive system. The diagnosis mostly depends on radiological examination. Surgery may be effective, but complications may occur.

Keywords: Zinner syndrome (ZS); symptom; diagnostic examinations; treatment; surgery; complication

Submitted Oct 10, 2020. Accepted for publication Dec 14, 2020.

doi: 10.21037/apm-20-1997

View this article at: <http://dx.doi.org/10.21037/apm-20-1997>

Introduction

Zinner syndrome (ZS) is a very rare anomaly, which is associated with an embryologic abnormality that develops in the distal portion of the mesonephric or Wolffian duct between the 4th and 13th gestational week. Incomplete migration of the ureteric bud from the proximal portion of the mesonephric duct may also result in a failure to meet with the metanephros. This failure interferes with the ureteric bud's key role in differentiation of the metanephric

blastema, and ipsilateral renal agenesis will occur, in addition to atresia of the ejaculatory duct. This leads to unsatisfactory drainage, subsequent distension of the seminal vesicle, and creation of a cystic structure (1).

The condition was first described by Zinner in 1914 (*Figure 1*). The incidence of ZS is very low. In one study of children in Taipei (2), the incidence of seminal vesicles with ipsilateral renal dysplasia was 0.0046% (13/28,000). The clinical symptoms associated with this condition, which are nonspecific, include frequent dysuria,

Figure 1 In a 30-year-old man with lower abdominal pain, CT presented the right renal agenesis (A) and posterior cystic lesions of the bladder (B,C, arrowheads). MRI showed the cystic lesions was attached to the right seminal vesicle and present as hyperintense in T1WI sequence and fat-suppression sequence (D,E) (short arrows) compared with bladder (stars) and fat (arrowheads). Operative and pathological findings as hemospermia in the seminal vesicle cyst. CT, computed tomography; MRI, magnetic resonance imaging.

epididymitis, perineal discomfort, and post-ejaculation pain. The severity of symptoms may correlate with the size of the seminal vesicle cyst (3). van den Ouden *et al.* (1) conducted a pooled analysis of patients with ZS in 1998. The study provided important reference information for clinicians and researchers; however, as times change and more cases are reported, an updated review of ZS cases may be necessary. We therefore performed a comprehensive review, which included case reports and case series published during the period from 1999 to 2020. We present our article in accordance with the PRISMA reporting checklist (available at <http://dx.doi.org/10.21037/apm-20-1997>).

Methods

A literature search was performed using the PubMed/Medline, Embase, and Scopus databases and the language was no restrictions. Two physicians with literature retrieval experience performed the search using the following keywords: zinner syndrome; zinnens syndrome; seminal vesicle cyst; ipsilateral renal agenesis; ipsilateral renal hypoplasia. Studies published during the period from January 1st, 1999 to May 1st, 2020 were included in the study. The data search was limited to case reports and

case series. In order to identify other potentially eligible publications, the references from the articles identified initially were manually reviewed. The diagnosis of ZS was considered in cases with of seminal vesicle cyst formation combined with ipsilateral renal agenesis.

The case reports were analyzed with regard to the data provided on patient age, clinical symptoms, site of pathogenesis, urogenital function, results on diagnostic examinations, treatment options, duration of follow-up, prognosis, complications, and comorbidities. The research is being reported in line with the attached PRISMA reporting checklist.

Statistical analysis

Descriptive statistics were used to summarize the variables in the case reports and case series.

Results

In total, 160 case reports and case series (Table S1) were obtained from the PubMed/Medline, Embase, and Scopus databases (Figure 2). All these cases met the proposed diagnostic criteria for ZS, and 214 patients were ultimately included.

Figure 2 Flow-chart depicting the screening process.

Patient characteristics at baseline

Twenty-one patients were removed due to a lack of data (4,5). The age of patients included in the study ranged from newborn (6) to 76 years (7) (mean age: 29.35 years) (*Table 1*). Of 214 patients, 173 (80.8%) had clinical symptoms, while the remaining cases were detected incidentally without symptoms (163%, 35/214), or had symptoms that were not described in detail (3.3%, 7/214). The symptoms described in the included studies were diverse, and most patients presented with multiple symptoms. Most patients presented with urinary symptoms, including: urinary frequency (24.2%, 42/173); dysuria (26.0%, 45/173); urinary urgency (13.9%, 24/173); micturition pain (8.2%, 14/173); hematuria (7.5%, 13/173); nocturia (1.7%, 3/173); urinary incontinence (1.2%, 2/173). The other common symptoms reported included: local pain, including perineal pain (20.2%, 35/173); abdominal pain (14.5%, 25/173); pelvic pain (6.9%, 12/173); suprapubic pain (4.0%, 7/173); backache (3.5%, 6/173); inguinodynia (2.9%, 5/173); testicular pain (1.7%, 3/173); scrotal pain

(1.2%, 2/173); coxalgia (0.6%, 1/173). Another category of symptoms in this patient population comprised abnormalities of reproductive function, including: infertility (13.9%, 24/173); ejaculation-related pain (11.0%, 19/173); hematospermia (8.7%, 15/173); testicular swelling (3.5%, 6/173); reduced or negligible semen volume (2.9%, 5/173); weak ejaculation (2.9%, 5/173); sexual discomfort (2.3%, 4/173); erectile dysfunction (1.2%, 2/173). Other symptoms reported included: constipation (4.0%, 7/173), fever (3.5%, 6/173), pain on defecation (1.2%, 2/173), loss of weight (1.2%, 2/173), nausea and vomiting (1.2%, 2/173), and so on. All reported symptoms are summarized in *Table 2* and *Figure 3*. In cases with seminal vesicle cyst enlargement and compression of the bladder, the most significant symptoms may be dysuria, perineal pain, urinary frequency, and urinary urgency. In 110 patients (51.4%), the seminal vesicle cyst and renal agenesis were present on the patient's right side. In 104 patients (48.6%), the seminal vesicle cyst and renal agenesis were present on the patient's left side. Thus, the patient's right side was more commonly affected, but the R:L ratio was approximately 1:1.

Diagnostic examinations

Table 3 summarizes the results of diagnostic examination for the 214 patients included in the study. Most patients underwent more than two diagnostic examinations. The techniques used most commonly for diagnosis were: magnetic resonance imaging (MRI) (145/214, 67.8%),

ultrasonography (139/214, 65%), computed tomography (CT) (130/214, 60.7%), urography (41/214, 19.2%), and digital rectal examination (DRE) (40/214, 18.7%).

Although radiology was most commonly used for the diagnostic examination, cystoscopy (30/214, 14.0%) and transrectal ultrasonography (TRUS) (29/214, 13.6%) were used to obtain additional information. Diagnostic methods such as retrograde pyelography (7/214, 3.3%) (8-14), radioactive nephrography (5/214, 2.3%) (9,15-18), positron emission tomography-computed tomography (PET-CT) (2/214, 0.9%) (19,20), and digital subtraction angiography (DSA) (2/214, 0.9%) (21,22) were effective in special situations.

Effects on urinary and reproductive function

When a seminal vesicle cyst compresses vital structures, maximum flow rate, as measured by uroflowmetry, may be attenuated. Uroflowmetry was performed in 10 cases. The maximum flow rate was 3.1 (18) to 16 (23) m/s, with median and standard deviation of 10.4 and 4.2 m/s, respectively. The normal reference range for maximum flow rate is ≥ 15 m/s.

Ejaculate volume was measured in 12 patients. Values

Table 1 Distribution of age (in decades) at the time of presentation in 193 ZS patients

Decade (year)	N	%
0–10	23	11.9
11–20	32	16.6
21–30	64	33.1
31–40	28	14.5
41–50	23	11.9
51–60	10	5.2
61–70	12	6.2
71–80	1	0.5

ZS, Zinner syndrome.

Table 2 Summary of symptoms in 173 ZS patients

Urinary symptoms		Local pain		Abnormal of reproductive function		Other symptoms	
Symptoms	Number	Symptoms	Number	Symptoms	Number	Symptoms	Number
Frequency	42	Perineal pain	35	Infertility	24	Constipation	7
Dysuria	45	Abdominal pain	25	Ejaculation pain	19	Fever	6
Urgent	24	Pelvic pain	12	Hematospermia	15	Defecation pain	2
Micturition pain	14	Suprapubic pain	7	Swelling of testicles	6	Loss of weight	2
Hematuria	13	Backache	6	Ejaculation weakness	5	Nausea and vomiting	2
Nocturia	3	Inguinodynia	5	Less or no semen	5	Diarrhea	1
Urinary Incontinence	2	Testicular pain	3	Sexual discomfort	4	Gastrointestinal uncomfortable	1
		Scrotal pain	2	Erectile dysfunction	2	Fatigued	1
		Coxalgia	1			Multiple	1
						Fremitus	1
						Abdominal distension	1
						Night sweats	1
						Malaise	1

ZS, Zinner syndrome.

Figure 3 The frequency of various symptoms in 173 ZS patients. ZS, Zinner syndrome.

ranged from 0.1 (24) to 2.2 (25) mL, with median and standard deviation of 1 and 0.6, respectively. The normal reference range for ejaculate volume is 2–6 mL. Details relating to sperm count were available for 7 patients. Among the patients included in this study, sperm count ranged from 160,000 (26) to 11,000,000 (27), with median and standard deviation of 1,200,000 and 5,200,000, respectively. The normal reference range for sperm count is >20,000,000. Data on sperm motility were available for 7 patients. The percentage of non-motile sperm in a given patient ranged from 59.5% (28) to 100% (24), with median and standard deviation of 90% and 18.35%, respectively. In an additional 12 cases, patients had been reported to have azoospermia and oligospermia, but details were not provided.

Treatment and management

Some of the cases included in this study did not describe

treatment. Details related to treatment and management were available for 193 patients (*Table 4*). Surgery, which was the most common approach to treatment, was performed in 127 cases (65.8%, 127/193), including 39 patients who underwent open surgery, 23 patients who underwent laparoscopic surgery, 23 patients who underwent robot-assisted laparoscopic vesiculectomy, five patients who underwent seminal vesiculoscopy, four patients who underwent transurethral resection of the ejaculatory duct, one patient who underwent ultrasound-guided transrectal biopsy, one patient who underwent interventional surgery, one patient who underwent cystoscopy and aspiration of the cyst, one patient who underwent transurethral unroofing of the cyst, and 29 patients who underwent surgery, with no details provided. In one case, death was caused by the metastasis of a papillary adenocarcinoma in the seminal vesicle after radical cystoprostatectomy (24). In another case, death was caused by multiple organ failure related to renal-type clear cell carcinoma in the prostate after surgery (29).

Table 3 Summary of diagnostic examination results for 214 ZS patients

Diagnostic examinations	N	The percentage of choice in 214 patients (%)
MRI	145	67.8
Trans-abdominal ultrasound	139	65.0
CT	130	60.7
Urography	41	19.2
DRE	40	18.7
cystoscopy	30	14.0
TRUS	29	13.6
Retrograde pyelography	7	3.3
Radioactive nephrogram	3	1.4
PET-CT	2	0.9
99mTc-DMSA	2	0.9
DSA	2	0.9

ZS, Zinner syndrome; MRI, magnetic resonance imaging; CT, computed tomography; DRE, digital rectal examination; TRUS, transrectal ultrasonography; PET-CT, positron emission tomography-computed tomography; DSA, digital subtraction angiography.

The other patients who underwent surgery did not experience symptom recurrence or other complications.

With regard to invasive procedures, 19 patients (19/193, 9.8%) underwent aspiration and drainage; 18 patients underwent transrectal aspiration of the cyst, and one patient underwent percutaneous drainage under ultrasound guidance. Nine patients (9/19, 47.4%) had cyst recurrence requiring surgical treatment, and one patient died of sepsis caused by an infection of the seminal vesicle cyst (30).

Forty-seven patients elected to receive conservative treatment and follow-up because of a lack of symptoms, the presence of only minor symptoms, or personal reasons (24.2%, 47/193). The range of follow-up time ranged from 3 months to 15 years. During follow-up, an increase in seminal vesicle cyst size necessitated surgery in 2 cases (31,32). Vasectomy was performed in one infant for recurrent epididymitis and orchitis. Nephrectomy was performed in three infants for unilateral multicystic dysplastic kidney. Kidney transplantation was performed in three infants for the treatment of bilateral dysplastic kidney (6).

Complications and comorbidities

In most cases, ZS is associated with a good prognosis. However, among the cases included in this study, 52 patients had complications or comorbidities, and three died (*Table 5, Figure 4*). According to the definitions of complications and comorbidities, as well as reviews of the literature and discussions with urologists and radiologists, we treated urogenital infection and inflammation as complications and treated tumors, developmental deformities, and other diseases as comorbidities. The urogenital infection and inflammation in collection cases including urinary infection (13 patients), epididymitis (6 patients) (6,33-37), prostatitis (2 patients) (38,39), orchitis (2 patients) (6,13), bladder infection (1 patient) (6), and sepsis caused by an infection of the seminal vesicle cysts, resulting in death (1 patient) (30). The most common comorbidities were the development of tumors, including adenoma or adenocarcinoma (40-45) arising from the seminal vesicle cyst (6 patients), paraganglioma affecting the bladder in 1 patient and affecting paranephrotic tissue in another patient (46,47), squamous cell carcinoma arising from the seminal vesicle (1 patient) (48), non-hodgkin lymphoma (1 patient) (19), and renal-type clear cell carcinoma affecting the prostate (1 patient) (43). Two patients with adenocarcinoma associated with a seminal vesicle cyst died because of metastasis and multiple organ failure. ZS may also be combined with other developmental deformities, including cryptorchidism (2 patients) (49,50), anorectal malformation (2 patients), contralateral dysplastic kidney (2 patients), ventricle septum defect (2 patients) (6), situs inversus viscerum (1 patient) (51), mitral insufficiency (1 patient) (52), and multiple severe developmental deformities (1 patient) (53). Comorbidities caused by unrelated illness or syndromes were relatively rare. In this study, 3 patients had complications related to Kallmann syndrome (21,54-56), 1 patient had complications related to thalassemia, and 1 patient had complications related to Fabry disease (57).

Discussion

ZS is a rare condition. The constellation of associated congenital findings includes ipsilateral renal hypoplasia, cystic seminal vesicles, and ejaculator duct obstruction. It occurs with anomalous growth of the mesonephric or

Table 4 Summary of management in 193 ZS cases

Management measures	N	%	Prognosis
Open operation	39	20.2	1 patient dead
Surgical operation no details	29	15.0	1 patient dead
Laparoscopic surgery	23	11.9	Favorable
Robotic-assisted laparoscopic vesiculectomy	23	11.9	Favorable
Seminal vesiculoscopy	5	2.6	Favorable
Transurethral resection of the ejaculatory duct	4	2.1	Favorable
Ultrasound-guided transrectal biopsy	1	0.5	Favorable
Interventional operation	1	0.5	Favorable
Cystoscope and aspiration of the cyst	1	0.5	Favorable
Transurethral unroofing of the cyst	1	0.5	Favorable
Aspiration and drainage	19	9.8	9 patients recurrence & 1 patient dead
Follow up	47	24.3	9 patients (included 7 infants) need surgery
Total	193	100.0	

ZS, Zinner syndrome.

Wolffian duct between 4 and 13 weeks of gestation and negatively impacts embryological development of the kidney and ejaculatory duct.

A pooled analysis was published by van den Ouden *et al.* in 1998. We sought to provide an updated review to help clinicians. Most patients with ZS present during the second or third decade of life. Compared with the study published by van den Ouden *et al.*, our study had a greater sample size. We found that there was no obvious difference in the incidence of left-sided *vs.* right-sided disease. This finding differs from those reported by van den Ouden *et al.* The most common symptoms among our patient population were dysuria, perineal pain, urinary frequency, and urinary urgency. These symptoms may be associated with an increase in cyst size. On the other hand, some patients had impaired reproductive function, including 12 patients presented with low-volume ejaculation, seven patients had low sperm motility, and 19 patients presented with azoospermia and oligospermia. Decreases in sperm count may be related to obliteration of the ejaculatory duct, the hemorrhage of seminal vesicle cysts, or secondary inflammation of the genitourinary system.

We also found that the choice of diagnostic examination had important effects on the selection of radiological techniques. In 1998, DRE and intravenous urography were commonly used for the diagnosis of ZS. In contrast,

the studies included in this review used MRI, trans-abdominal ultrasound, and CT because of these techniques' noninvasive nature and ease of use. MRI was recommended by radiologists to differentiate between lesions and surrounding normal tissue as well as the effects of non-ionizing radiation. Recent developments in the field of MRI have led to improved field intensity and new sequences, which provide more information about the condition of underdeveloped kidneys and ureters, cyst contents, and surrounding tissue or organs affected by complications. This information is helpful for guiding the clinical decision. In this review, seminal vesicle cyst and ipsilateral renal agenesis were first detected by trans-abdominal ultrasound because of its convenience for screening. CT is an important supplement for patients with contraindications for MRI. The information obtained via CT and urography can supplement other information such as the extent of calcification and urinary function.

In the studies included in this review, surgery (open or laparoscopic) was the most effective treatment and postoperative recurrence was rare. As new methods of surgery that were not mentioned in the study by van den Ouden *et al.*, endoscopy and robot-assisted technology have become increasingly popular among physicians performing seminal vesiculectomy since 2002 (58) and 2007 (59), respectively. Compared with open seminal vesiculectomy,

Table 5 Complications and comorbidities in 52 ZS patients

Complications and comorbidities	N	The percentage in 214 patients (%)
Tumors	11	
Adenocarcinoma (seminal vesicle)	5	2.34
Adenoma	1	0.47
Paraganglioma (bladder and aranephros)	2	0.93
Squamous cell carcinoma (seminal vesicle)	1	0.47
Non-Hodgkin lymphoma (lymphoid tissue)	1	0.47
Renal-type clear cell carcinoma (prostate)	1	0.47
Developmental deformities	10	
Situs inversus viscerum	1	0.47
Mitral insufficiency	1	0.47
Cryptorchidism	2	0.93
Anorectal malformation	2	0.93
Contralateral dysplastic kidney	2	0.93
Ventricle septum defect	2	0.93
Multiple developmental deformity	1	0.47
Infections or inflammations	25	
Urinary infection	13	6.07
Bladder infection	1	0.47
Epididymitis	6	2.80
Prostatitis	2	0.93
Orchitis	2	0.93
Sepsis	1	0.47
Other illness or syndromes	5	
Kallmann syndrome	3	1.40
Thalassemia	1	0.47
Fabry disease	1	0.47
Total number of patients	52	23.83

ZS, Zinner syndrome.

laparoscopic and robot-assisted seminal vesiculectomy are safer, less invasive, accompanied by less bleeding, and associated with quicker recover. Among the 46 patients included in this review who underwent laparoscopic (23 patients) or robot-assisted seminal vesiculectomy (23 patients), prognosis was favorable, with no postoperative complications reported. However, our sample size was

insufficient to demonstrate the safety of surgical treatment for ZS. A large-scale of meta-analysis (60) that included more than 200,000 patients indicated that open retropubic radical prostatectomy (ORP), laparoscopic radical prostatectomy (LRP), and robot-assisted laparoscopic radical prostatectomy (RALP) were associated with high risk for surgical complications leading to erectile dysfunction (ORP: 42.7–61.8%; LRP: 45–71.1%; RALP: 23.4–68.2%) and urinary incontinence (ORP: 15.4–28%; LRP: 7.1–42.8%; RALP: 7.7–20.7%). Although the target organs are different, the surgical methods, surgical approaches, and surrounding tissues are similar between prostatectomy and seminal vesicle resection, seminal vesicle resection still carries risk for surgical complications. Postoperative recurrence was not found in the van den Ouden *et al.* (1) report or our population, but that does not mean that it is impossible. Additional cases and longer follow-up time are necessary.

Another option in such cases is aspiration and drainage. van den Ouden *et al.* reported that the rate of success after aspiration was 30%, and 47.4% of patients included in the present review who underwent aspiration had symptom recurrence. As a variety of invasive treatment, aspiration was unable to decrease the risk for seminal vesicle cyst recurrence, so it is not valuable as a management strategy. Asymptomatic patients should be followed with observation, but when symptoms appear, clinical intervention is needed.

The complications and comorbidities experienced by ZS patients deserve attention but have been neglected in the past. In this review, three patients died because of complications or comorbidities. Secondary inflammation and ZS-related are the frequent complications in this patient population. These complications present with symptoms, so they should not easily be overlooked. However, tumor as comorbidity may not present with any specific symptoms in the early stage. In our review, tumors occurred in 10 patients (4.4%), and two patients died because of multiple metastases or organ failure. Patients with ZS and tumors arising from the urogenital system [including adenocarcinoma of the seminal vesicle (5, 2.34%), clear cell renal carcinoma of the prostate (1, 0.47%), paraganglioma (2, 0.93%) and squamous cell carcinoma of the seminal vesicle (1, 0.47%)] are rare conditions in ordinary people. According to the relatively high incidence of disease in this ZS population, although the exact reason remains unknown, ZS may be a positive stimulating factor for urogenital system tumors. Other developmental abnormalities or diseases can occur, but they are relatively rare. Remarkably,

Figure 4 The frequency of complications and comorbidities in 52 ZS patients. ZS, Zinner syndrome.

three ZS patients included in this review had Kallmann syndrome, which was more common than cryptorchidism and contralateral dysplastic kidney. We consider that ZS and Kallmann syndrome may have some kind of relationship, but the cases and data were not sufficient for investigation.

This review has several limitations. First, it was not possible to classify patients by ethnicity because the information was lacking from many studies. Only 13 patients could be classified according to ethnicity: 7 Caucasian males; 2 Saudi Arabian males; 1 Indian male; 1 Hispanic male; 1 Moroccan male; 1 Italian male. Second, it was not possible to classify seminal vesicle lesions based on wall changes because the information was lacking from many studies. Third, there may have been some publication bias, resulting in overestimations of the proportions of symptoms, complications, and comorbidities. Fourth, a majority of cases lack relevant information such as uroflowmetry, semen ejaculate volume, sperm count, and sperm motility, so it difficult to provide convincing data on the effect of ZS on male reproductive function. Finally, some of the studies and conference papers identified by our initial literature

search provided only abstracts. We had to exclude these publications from our review. In conclusion, the clinical symptoms associated with ZS syndrome are varied, but the most common symptoms are enlargement of seminal vesicle cysts and compression of the bladder, ureters, and reproductive system, as well as associated secondary lesions. The diagnosis mostly depends on the results of radiological examinations. Complications and comorbidities should be identified, with treatment tailored accordingly.

Acknowledgments

Funding: None.

Footnote

Reporting Checklist: The authors have completed the PRISMA reporting checklist. Available at <http://dx.doi.org/10.21037/apm-20-1997>

Conflicts of Interest: All authors have completed the ICMJE

uniform disclosure form (available at <http://dx.doi.org/10.21037/apm-20-1997>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- van den Ouden D, Blom JH, Bangma C, et al. Diagnosis and management of seminal vesicle cysts associated with ipsilateral renal agenesis: a pooled analysis of 52 cases. *Eur Urol* 1998;33:433-40.
- Sheih CP, Hung CS, Wei CF, et al. Cystic dilatations within the pelvis in patients with ipsilateral renal agenesis or dysplasia. *J Urol* 1990;144:324-7.
- Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* 2020;382:727-33.
- Pace G, Galatioto GP, Gualà L, et al. Ejaculatory duct obstruction caused by a right giant seminal vesicle with an ipsilateral upper urinary tract agenesis: an embryologic malformation. *Fertil Steril* 2008;89:390-4.
- Jiang XS, Wang HJ, Lin JH, et al. Zinner's syndrome: clinical features and imaging diagnosis. *Asian J Androl* 2018;20:316-7.
- Schukfeh N, Kuebler JF, Schirg E, et al. Dysplastic kidney and not renal agenesis is the commonly associated anomaly in infants with seminal vesicle cyst. *BJU Int* 2009;103:816-9.
- Fujita M, Goto T, Saiki S. Seminal vesicle cyst associated with ipsilateral renal dysplasia and vena cava malformation: a case report. *Hinyokika Kiyo* 2012;58:511-4.
- Burns JA, Cooper CS, Austin JC. Cystic dysplasia of the testis associated with ipsilateral renal agenesis and contralateral crossed ectopia. *Urology* 2002;60:344.
- Sukumar S, Khanna V, Nair B, et al. Adult presentation of congenital ectopic vas deferens insertion into ureter with unilateral renal agenesis. *Surg Radiol Anat* 2010;32:9-10.
- Nayak B, Abrol N, Kumar R. Radiological seminal vesicle stones may actually be in the ureter. *Indian J Urol* 2013;29:75-6.
- Haddock P, Wagner JR. Seminal vesicle cyst with ipsilateral renal agenesis and ectopic ureter (Zinner syndrome). *Urology* 2015;85:e41-2.
- Sundar R, Sundar G. Zinner syndrome: an uncommon cause of painful ejaculation. *BMJ Case Rep* 2015;2015:bcr2014207618.
- Cascini V, Di Renzo D, Guerriero V, et al. Zinner syndrome in pediatric age: issues in the diagnosis and treatment of a rare malformation complex. *Front Pediatr* 2019;7:129.
- Maré A, Liddell H, Chan HF. Ectopic megaureter inserting into seminal vesicle cyst with fistula to bladder. *ANZ J Surg* 2017;87:E90-1.
- Chuang KL, Lai WI, Chiang YJ. Giant seminal vesicle cyst resembling megaureter with hydronephrosis. *Arch Androl* 2005;51:367-9.
- Ulu EMK, Başaran C, Dönmez FY, et al. Mullerian duct cyst with ipsilateral renal agenesis: MRI findings. *Marmara Medical Journal* 2009;22:64-7.
- Jaiswal A, Francis SK, Tuli A, et al. Zinner syndrome: rare cause of urinary symptoms in young adult male. *Indian J Urol* 2016.
- Maehana T, Fukuta F, Kobayashi K, et al. Laparoscopic surgery for seminal vesicle cysts and ureterocele with urination disorder: a case report of Zinner syndrome. *J Endourol Case Rep* 2018;4:35-8.
- Kuo J, Foster C, Shelton DK. Zinner's syndrome. *World J Nucl Med* 2011;10:20-2.
- Harbo FS, Larsen LB. Seminal vesicle cyst with ipsilateral renal agenesis. *Ugeskr Laeger* 2015;177:104-5.
- Savica V, Santoro D, Monardo P, et al. Seminal vesicle cysts with unilateral renal agenesis and contralateral ureteral stenosis in a β -thalassemic patient: an unknown association by incomplete development of the mesonephric duct. *Urol Int* 2007;79:367-70.
- Royston E, Walker M, Ching B, et al. Postcoital hemorrhage of a recurrent seminal vesicle cyst requiring embolization. *Urol Case Rep* 2014;2:171-2.
- Kiremit MC, Acar O, Sag AA, et al. Minimally invasive management of Zinner's syndrome with same-session robot-assisted seminal vesiculectomy and ipsilateral nephroureterectomy using a single geometry of trocars. *J Endourol Case Rep* 2018;4:186-9.

24. Kawahara T, Nishiyama H, Okubo K, et al. Right ectopic ureter with ipsilateral renal agenesis presenting with infertility: a case report. *Hinyokika Kyo* 2004;50:435-8.
25. Andrade-Rocha FT. Semen analysis in an infertile man with seminal vesicles cysts associated with ipsilateral renal agenesis. *Int Urol Nephrol* 2006;38:101-3.
26. Calahorra Fernández FJ, García Cardoso JV, Montero Rubio R, et al. Quiste gigante de vesícula seminal asociado con agenesia renal ipsilateral. *Actas Urol Esp* 2002;26:218-23.
27. Mehra S, Ranjan R, Garga UC. Zinner syndrome-a rare developmental anomaly of the mesonephric duct diagnosed on magnetic resonance imaging. *Radiol Case Rep* 2016;11:313-7.
28. Kao CC, Wu CJ, Sun GH, et al. Congenital seminal vesicle cyst associated with ipsilateral renal agenesis mimicking bladder outlet obstruction: a case report and review of the literature. *Kaohsiung J Med Sci* 2010;26:30-4.
29. Sato Y, Kataoka M, Hata J, et al. Renal-type clear cell carcinoma occurring in the prostate with Zinner syndrome. *Urol Case Rep* 2016;5:9-12.
30. Chung SD, Sun HD, Chiu CY. Lethal infected seminal vesicle cyst. *Int J Infect Dis* 2009;13:e197-8.
31. te Dorsthorst MJ, Prins-van Uden M, Niekel R, et al. Case report. Zinner-syndroom: een zeldzame oorzaak van agenesie van de nier. *Tijdschrift voor Urologie* 2020;10:9-12.
32. Alhajeri F, Salem S, Harraz A, et al. Scientific Program of 37th World Congress of Endourology Program Book. *J Endourol* 2019;33.
33. Giglio M, Medica M, Germinale F, et al. Renal dysplasia associated with ureteral ectopia and ipsilateral seminal vesicle cyst. *Int J Urol* 2002;9:63-6.
34. Liatsikos EN, Lee B, Filos KS, et al. Congenital seminal vesicle cyst and coexisting renal agenesis: laparoscopic approach. *Urology* 2004;63:584-6.
35. Berg F, Ohlmann C, Gossmann A. MR tomographic diagnosis of a congenital seminal cyst in ipsilateral renal agenesis with ureterocele as rare differential diagnosis of prostatic tumor. *Rofo* 2005;177:1308-9.
36. Heller MT, Hartman M, McGreevy B. Seminal vesicle cysts causing pelvic pain: importance of computed tomography. *Am J Emerg Med* 2012;30:2087.e1-6.
37. Herrero Blanco E, Campos-Juanatey F, Lastra García-Barón P, et al. Zinner's syndrome in a polytrauma patient. *Arch Esp Urol* 2019;72:608-11.
38. McAdams S, Schomburg J, Sweet RM. Zinner's syndrome. Like sitting on a tennis ball for 20 years. *Minn Med* 2014;97:40-1.
39. Sigüenza González R, Sánchez Lite I. Seminal vesicle cyst: An isolated finding? *Radiologia* 2018;60:446-8.
40. Kluckert JT, Zaunbauer W, Diener PA. Congenital seminal vesicle cyst with an intracystic papillary adenoma associated with ipsilateral renal agenesis. *Radiologe* 2002;42:837-9.
41. Kondo N, Shiono Y, Yoshino Y, et al. Papillary adenocarcinoma in a seminal vesicle cyst associated with contralateral renal agenesis: a case report. *Hinyokika Kyo* 2007;53:175-8.
42. Lee BH, Seo JW, Han YH, et al. Primary mucinous adenocarcinoma of a seminal vesicle cyst associated with ectopic ureter and ipsilateral renal agenesis: a case report. *Korean J Radiol* 2007;8:258-61.
43. Satyanarayana R, Katkooi D, Young E, et al. Robotic assisted excision of a giant seminal vesicle cyst and atretic ureter. *J Endourol* 2013;27:A443.
44. Deptala A, Romanowicz A, Czerw A, et al. Congenital anomalies of the genitourinary system can help in diagnosis of the primary site of metastatic cancer: a case report and a review of the literature. *Onco Targets Ther* 2016;9:4435-40.
45. Bhat A, Banerjee I, Kryvenko ON, et al. Primary seminal vesicle adenocarcinoma: a lethal yet cryptic malignancy with review of literature. *BMJ Case Rep* 2019;12:e232994.
46. Valsangkar RS, Goyal NK, Bajania SP, et al. Bladder paraganglioma with renal agenesis: A possible new association and its implications in the light of rearranged in transfection gene genetics. *Urol Ann* 2015;7:410-3.
47. Ali MS, Dubey D, Dias B. Robotic Assisted excision of extra adrenal Paraganglioma presenting as left seminal vesicle mass. *Indian J Urol* 2018;34:37.
48. Kim YH, Moon KC. Zinner Syndrome patient with squamous cell carcinoma arising from squamous metaplasia of the seminal vesicle: a case report. *Virchows Archiv* 2014;465:S165.
49. Yu YD, Hong YK. A case of cryptorchidism with ipsilateral congenital unilateral absence of the vas deferens and contralateral renal agenesis. *Case Rep Urol* 2016;2016:2379793.
50. Naval-Baudin P, Carreño García E, Sanchez Marquez A, et al. Multicystic seminal vesicle with ipsilateral renal agenesis: two cases of Zinner syndrome. *Scand J Urol* 2017;51:81-4.
51. Fiaschetti V, Greco L, Giuric, et al. Zinner syndrome diagnosed by magnetic resonance imaging and computed tomography: role of imaging to identify and evaluate the

- uncommon variation in development of the male genital tract. *Radiol Case Rep* 2016;12:54-8.
52. Kowalczyk K, Baka-Ostrowska M, Felberg K, et al. Unilateral renal dysplasia associated with ectopic ureter opening into ipsilateral ejaculatory duct. *Cent European J Urol* 2009;62:181-2.
 53. Lilje C, Finger LJ, Ascuitto RJ. Complete unilateral leg duplication with ipsilateral renal agenesis. *Acta Paediatr* 2007;96:464-5.
 54. Sridhar AN, Zacharakis E, Dudderidge T, et al. Robot-assisted management of Zinner's syndrome: report of seminal vesicle sparing technique and review of literature. *J Robot Surg* 2014;8:185-7.
 55. Domínguez A, Abascal-Junquera JM, Muñoz-Rodríguez J, et al. Síndrome de Zinner en paciente afecto de síndrome de Kallmann: rara asociación de 2 malformaciones embrionarias. *Revista Internacional de Andrología* 2016;14:144-7.
 56. Leen A, Dirk V. Zinner's syndrome in a patient with X-linked Kallmann syndrome: case report. *Acta Clin Belg* 2016;71:6.
 57. Sousa C, Teixeira IP, Ferreira SH, et al. Zinner's syndrome: case report of a rare maldevelopment in the male genitourinary tract. *Journal of Urological Surgery* 2019;6:65.
 58. Buogo G, Rodrigues H, Rodrigues P. Laparoscopic removal of seminal vesicle cyst with ectopic ureteral insertion and renal remnant. *Int Braz J Urol* 2002;28:335-7.
 59. Moore CD, Erhard MJ, Dahm P. Robot-assisted excision of seminal vesicle cyst associated with ipsilateral renal agenesis. *J Endourol* 2007;21:776-9.
 60. Basiri A, de la Rosette JJ, Tabatabaei S, et al. Comparison of retropubic, laparoscopic and robotic radical prostatectomy: who is the winner? *World J Urol* 2018;36:609-21.

Cite this article as: Liu T, Li X, Huang L, Li H, Cai K, Jiang J, Chen N, Zhang W, Tang J, Zhang M, Zhao D, Ye J, Zhang C, He T, Liu B, Ye Y, Chen J. Zinner syndrome: an updated pooled analysis based on 214 cases from 1999 to 2020: systematic review. *Ann Palliat Med* 2021;10(2):2271-2282. doi: 10.21037/apm-20-1997

Table S1 Distribution, clinical symptoms and ages features of Zinner syndrome (ZS) from 160 case report/case series

No.	Author	Number & distribution (R = right, L= left)	Age (y = year, m = month)	Symptoms and complains
1	Landry JL (61)	1R	24 y	Perineal pain
2	Steffens J (62)	1R	10 m	Asymptom
3	Livingston L (63)	3L & 2R	17–60 y	Dysuresia [4], perineal pain, urgent, frequency [2], infertility [2], nocturia, hematuresis, hematospermia, abdomen pain & ejaculation pain
4	Gorrea M (64)	1R	37 y	Hematuresis
5	Eberli D (65)	1R	9 y	Swelling of Testicles
6	Burns JA (8)	1R	12 y	Swelling of Testicles
7	Buogo G (58)	1L	23 y	Inguinodynia & sexual discomfort
8	Manousakas T (66)	1L	30 y	Hematuresis, frequency & urgent
9	Radhia S (67)	1R	34 y	Dysuresia, frequency, urgent & pelvic pain
10	Takada S (68)	1L	39 y	Asymptom
11	Calahorra Fernández FJ (26)	1R	41 y	Hematospermia & infertility
12	Giglio M (33)	1L	45 y	Epididymitis
13	Roberts SG (69)	1R	49 y	Asymptom
14	Kluckert JT (40)	1L	67 y	Dysuresia
15	Hoschke B (70)	1L	14 y	Asymptom
16	Protogerou V (71)	1L	24 y	Frequency, urgent, backache,inguinodynia
17	Segawa N (72)	1R	29 y	Uroschesis & perineal pain
18	Akao J (73)	1R	29 y	Dysuresia
19	Narlawar RS (74)	1R	30 y	Dysuresia, frequency & ejaculation pain
20	Murphy JO (75)	1L & 1R	15y & 22y	Dysuresia [2] & suprapubic pain [1]
21	Basillote JB (76)	1R	7y	Dysuresia & suprapubic pain
22	Kilinç F (77)	1L	17 y	Dysuresia & perineal pain
23	Liatsikos EN (34)	1L	26 y	Dysuresia, frequency & urgent
24	Kawahara T (24)	1R	30 y	Infertility
25	Pascual Samaniego M (78)	1R	34 y	Inguinodynia
26	Madrid García FJ (79)	1L	42 y	Dysuresia & frequency
27	Berg F (35)	1R	38 y	Perineal pain & ejaculation pain
28	Chuang KL (15)	1R	53 y	Abdominal pain
29	Kosan M (80)	1L	20 y	Hematuresis
30	Gözen AS (81)	1R	22 y	Dysuresia, frequency & uracratia
31	Romero Selas E (82)	1R	26 y	Asymptom
32	Andrade-Rocha FT (25)	1R	39 y	Infertility
33	Pal D (83)	1L	51 y	Dysuresia
34	Cihan A (84)	1L	61 y	Asymptom
35	Chen HW (85)	4L & 2R	10–15 y	Asymptom [4], dysuresia, frequency & urgent [1]
36	Savica V (21)	1R	13 y	Coxalgia
37	Moore CD (59)	1R	16 y	Nausea, vomiting, abdominal pain & swelling of testicles
38	Kondo N (41)	1R	30 y	Hematospermia, perineal pain, micturition pain
39	Suárez Aliago B (86)	1L	33 y	Abdominal pain & fever
40	Lee BH (42)	1L	41 y	Hematuresis
41	Han P (87)	1L	41 y	Perineal pain & hematospermia
42	Carbone A (88)	1 & 1R	24 y & 43 y	Abdominal pain & infertility [1], no semen [1]
43	Pace G (4)	7R	32 y (mean)	Infertility [7], ejaculation weakness [5], dysuresia [4], ejaculation pain [4] & perineal pain [4]
44	Lilje C (53)	1R	5 m	Multiple developmental malformation
45	Labanaris AP (89)	1R	27 y	Sexual discomfort, frequency & urgent
46	Casey RG (90)	1R	32 y	Swelling of testicles & testicular pain
47	Štimac G (91)	1R	36 y	Inguinodynia, frequency & urgent
48	Ohgaki K (92)	1L	38 y	Perineal pain & micturition pain
49	Selli C (93)	1R	39 y	Perineal pain & dysuresia
50	Kowalczyk K (52)	1R	3 y	Swelling of testicles
51	Sotelo R (94)	1R	19 y	Micturition pain, frequency & urgent
52	Pereira BJ (95)	1R	22 y	Dysuresia, perineal pain & ejaculation pain
53	Ulu EMK (16)	1R	25 y	Asymptom
54	Chung SD (30)	1R	56 y	Fever, dysuresia& uroschesis
55	Seo IY (96)	4L	22 y, 26 y, 67 y & 68 y	Perineal pain [2] & hematospermia [2]
56	Schukfeh N (6)	5L & 4R	Newborn to 50 m	Asymptom [8], epididymitis and orchitis [1]
57	Ghonge NP (97)	1R	19 y	Dysuresia, frequency, perineal pain, scrotal pain & ejaculation pain
58	Sukumar S (9)	1R	20 y	Abdominal pain
59	Lim Christopher SH (98)	1R	23 y	Ejaculation pain
60	Viktorsdóttir MB (99)	1R	25 y	Perineal pain
61	Allaparthi S (100)	1R	34 y	No mention
62	Dombóvári P (101)	1L	34 y	Abdominal pain
63	Kao CC (28)	1L	43 y	Perineal pain & infertility
64	Ahallal Y (102)	1L	16 y	Micturition pain, frequency & urgent
65	Hampton LJ (103)	1L	35 y	Micturition pain, less or no semen & epididymitis
66	Kovac JR (104)	1L	36 y	Abdominal pain, astriction, nausea, vomiting & fever
67	Jang KD (105)	1L	49 y	Frequency, nocturia, & dysuresia
68	Kuo J (19)	1L	52 y	Fever, loss of weight & night sweats
69	Kalyanaraman B (106)	1L	20 y	Perineal pain, dysuresia & defecation pain
70	Devaraju S (107)	1R	28 y	Abdominal pain, dysuresia, frequency & suprapubic pain
71	Basal S (108)	1L	28 y	Perineal pain & dysuresia
72	El-Assmy A (109)	1R	34 y	Frequency & nocturia
73	Haghighi R (110)	1L	36 y	Perineal pain & hematospermia
74	Fujita M (7)	1R	76 y	Dysuresia
75	Heller MT (36)	2R	46 y, 27 y	Perineal pain [2]
76	Coşkun B (111)	1L	20 y	Uracratia, pelvic pain, sexual discomfort & hematospermia
77	Kardoust Parizi M (112)	1L	27 y	Hematospermia, ejaculation pain, infertility, micturition pain, frequency & urgent
78	Sridhar AN (54)	1R	36 y	Kallmann syndrome, hematuresis & hematospermia
79	Nayak B (10)	1R	44 y	Dysuresia, hematuresis & frequency
80	Altobelli E (113)	1L	66 y	Fever & fremitus
81	Silveri M (114)	1L	2 y	Abdominal pain & astriction
82	Batur A (115)	1L	22 y	Pelvic pain & hematospermia
83	Royston E (22)	1L	23 y	Ejaculation pain
84	Palmer WC (116)	1L	26 y	Fever and pelvic pain
85	Satyanarayana R (43)	1R	29 y	No mention
86	Kim YH (48)	1R	29 y	Less or no semen & hematuresis
87	Lee KH (117)	1L	46 y	Abdominal pain, hematuresis, dysuresia, micturition pain, frequency & urgent
88	Mcadams S (38)	1R	54 y	Dysuresia, frequency & pelvic pain
89	Shepherd G (118)	2L	18 m & 2 y	Asymptom [2]
90	Kanavaki A (119)	1R	4 y	Asymptom
91	Burak Özkan M (120)	1L	15 y	Pelvic pain
92	Haddock P (11)	1R	18 y	Ejaculation pain
93	Valsangkar RS (46)	1R	21 y	Hematuresis
94	Donato P (121)	1L	22 y	Abdominal pain & diarrhea
95	Sundar R (12)	1L	25 y	Frequency, pelvic pain & ejaculation pain
96	Alp B (122)	1L	28 y	Frequency & pelvic pain
97	Harbo FS (20)	1L	41 y	Swelling of testicles
98	Kim Y (123)	1R	41 y	Hematuresis
99	Pavan N (124)	1R	42 y	Asymptom
100	Juho YC (125)	1R	43 y	Gastrointestinal uncomfortable, less or no semen & frequency
101	García Asensio D (126)	1L	10 m	Asymptom
102	Domínguez A (55)	1R	20 y	Kallmann syndrome, testicular pain & erectile dysfunction
103	Leen A (56)	1L	22 y	Kallmann syndrome & testicular hypoplasia
104	Jaiswal A (17)	1R	25 y	Micturition pain, frequency & urgent
105	Mehra S (27)	1R	25 y	Dysuresia, frequency & infertility
106	Aghaways I (127)	1L	28 y	Abdominal pain
107	Hajji F (128)	1L	29 y	Dysuresia, urgent, abdominal pain, sexual discomfort & hematospermia
108	Dogan F (129)	1R	38 y	Hematospermia, dysuresia, astriction & pelvic pain
109	Shah S (130)	1R	39 y	Uroschesis & suprapubic pain
110	Bakloul F (131)	1R	43 y	Infertility
111	Gutierrez García MA (132)	1L	46 y	Pelvic pain & ejaculation pain
112	Deptala A (44)	1L	58 y	Frequency, backache, loss of weight & astriction
113	Maré A (133)	1R	62 y	No mention
114	Sato Y (29)	1L	64 y	Hematuresis
115	Yu YD (49)	1R	10 m	Asymptom & undescended testis
116	Slaoui A (134)	1L & 1R	35 y & 39 y	Hematuresis [1], backache [1] & no mention [1]
117	Canales-Casco N (135)	1L	17 y	Hematospermia
118	Prem G (136)	1L	19 y	Dysuresia, frequency, urgent & ejaculation pain
119	Chiapparrone G (137)	1L	20 y	Perineal pain & infertility
120	Cao JD (138)	1L	23 y	Perineal pain
121	Ahmed A (139)	1L	30 y	Backache & abdominal pain
122	Campora M (140)	1R	37 y	Asymptom
123	Khanduri S (141)	1R	40 y	Abdominal pain, perineal pain, ejaculation pain & infertility
124	Fiaschetti V (51)	1L	56 y	Perineal pain, dysuresia, frequency & less or no semen
125	Naval-Baudin P (50)	1L & 1R	26 y & 29 y	Testicular pain [1] & infertility [1]
126	Roth C (142)	1L	15 y	Perineal pain & frequency
127	Millman AL (143)	1R	21 y	Frequency, urgent & perineal pain
128	Maehana T (18)	1R	21 y	Dysuresia
129	Razdan S (144)	1L	21 y	Abdominal pain, pelvic pain & astriction
130	Florim S (145)	1R	21 y	Scrotal pain
131	Kord E (146)	1L	23 y	Micturition pain, frequency, urgent & perineal pain
132	Kiremit MC (23)	1L	23 y	Micturition pain, frequency, urgent & perineal pain
133	Siguenza Gonzalez R (39)	1R	23 y	No mention
134	Farooqui A (147)	1R	28 y	Malaise, fatigue, & myalgia
135	Brown M (148)	1L	31 y	Micturition pain, frequency & urgent
136	El Mortaji H (149)	1R	45 y	Backache
137	Ali MS (47)	1L	67 y	Micturition pain, frequency, urgent & ejaculation pain
138	Guan J (5)	6L & 8R	14–43 y	Hematospermia [1], infertility [2], abdominal pain [3], ventosity [1], frequency & dysuresia [7]
139	Bríosa F (150)	1R	14 y	Asymptom
140	Sousa Célia (57)	1R	17 y	No mention
141	Symeonidis EN (151)	1L	19 y	Asymptom
142	Kori R (152)	1R	19 y	Dysuresia
143	AlArifi M (153)	1L	20 y	Inguinodynia, testicular pain & ejaculation pain
144	Aslan Serdar (154)	1L	27 y	Pelvic pain
145	Cito G (155)	1L	27 y	Perineal pain & dysuresia
146	Bhat A (45)	1R	28 y	Perineal pain & abdominal pain
147	Bhukte S (156)	1L	28 y	Abdominal pain
148	Patil M (157)	1R	32 y	Infertility
149	Sahare P (158)	1R	45 y	Micturition pain, frequency, urgent & dysuresia
150	Alhajeri F (159)	1L	46 y	Testicular pain, suprapubic pain & uroschesis
151	Corongiu E (160)	1L	51 y	Perineal pain, astriction, infertility
152	Chibelean BC (161)	1L	61 y	Abdominal pain & astriction
153	Cascini V (13)	1L & 1R	1 m & 1 m	Asymptom
154	Herrero Blanco E (37)	1L & 1R	22 y, 29 y	Asymptom [2]
155	Bryson CF (162)	1R	24 y	Abdominal pain, suprapubic pain, testicular pain and frequency
156	Hergan B (163)	1L	22 y	Asymptom
157	Yadav SP (164)	1L	29 y	Suprapubic pain, uroschesis,
158	Tan Z (165)	2L & 4R	15–64 y	Micturition pain [1], Frequency [2], Urgent [2] Backache [1], Asymptom [2], hematochezia [1]
159	te Dorsthorst MJ (31)	2L& 1R	56 y, 61 y, 67 y	Asymptom [1], erectile dysfunction [1], abdominal pain [1], uroschesis [1]
160	Jatal S (166)	1L	28 y	Frequency, urgent & defecation pain

References

61. Landry JL, Dodat H, Pelizzo G, et al. Cystic dysplasia of the rete testis and ipsilateral kidney agenesis in children. *Arch Pediatr* 1999;6:416-20.
62. Steffens J, Oberschulte-Beckmann D, Siller V, et al. Ectopic refluxing ureter entering a seminal vesicle cyst associated with ipsilateral renal dysplasia. *World J Urol* 2000;18:232-4.
63. Livingston L, Larsen CR. Seminal vesicle cyst with ipsilateral renal agenesis. *AJR Am J Roentgenol* 2000;175:177-80.
64. Gorrea M, Lorente R, Roel J. Seminal vesicle cyst associated with ipsilateral renal agenesis and papillary carcinoma of the bladder. *Eur Radiol* 2001;11:2500-3.
65. Eberli D, Gretener H, Dommann-Scherrer C, et al. Cystic dysplasia of the testis: a very rare paediatric tumor of the testis. *Urol Int* 2002;69:1-6.
66. Manousakas T, Kyriakou G, Serafetinides E, et al. Partial vesiculectomy in an infertile man with seminal vesicle cyst, ipsilateral renal agenesis, and cryptorchidism. *Urology* 2002;59:602.
67. Radhia S, Samira F, Mounir T, et al. Kyste de la vésicule séminale associé à une agénésie rénale et à un uretère ectopique ipsilatéral. À propos d'un cas. *Annales d'Urologie* 2002;36:381-3.
68. Takada S, Noda Y, Oka T, et al. Seminal vesicle cyst associated with ipsilateral renal dysplasia: a case report. *Nishinihon Journal of Urology* 2002;64:436-9.
69. Roberts SG, Garcia Mediero JM, Segura JW, et al. Incidental pelvic mass identified during ultrasound-guided transrectal needle biopsy of the prostate. *Arch Esp Urol* 2002;55:466-8.
70. Hoschke B, May M, Seehafer M, et al. Outlet of a megaureter with aplastic kidney into a seminal vesicle cyst. Case report of laparoscopic intervention. *Urologe A* 2003;42:1092-6.
71. Protogerou V, Chorti M, Kostakopoulos A. Congenital dilation of seminal vesicle with agenesis of ipsilateral kidney: symptoms and treatment. *Int Urol Nephrol* 2003;35:69-71.
72. Segawa N, Gohji K, Masuda H, et al. A case of CA19-9-producing seminal vesicle cyst with ipsilateral renal agenesis. *Hinyokika Kiyo* 2003;49:277-80.
73. Akao J, Arikawa M, Aramaki K, et al. Seminal vesicle cyst associated with ipsilateral renal dysplasia: a case report. *Nishinihon Journal of Urology* 2003;65:505-8.
74. Narlawar RS, Hanchate V, Raut A, et al. Renal agenesis and seminal vesicle cyst. *J Ultrasound Med* 2003;22:225-8.
75. Murphy JO, Power RE, Akhtar M, et al. Magnetic resonance imaging in the diagnosis of seminal vesicle cysts and associated anomalies. *J Urol* 2003;170:2386.
76. Basillote JB, Shanberg AM, Woo D, et al. Laparoscopic excision of a seminal vesicle cyst in a child. *J Urol* 2004;171:369-71.
77. Kiliç F, Güvel S, Eğilmez T, et al. Seminal vesicle cyst, ectopic ureter opening into the cyst and ipsilateral renal agenesis. *Türk Uroloji Dergisi* 2004;30:369-71.
78. Pascual Samaniego M, Egea Camacho J, Cortiñas González JR, et al. Agenesia renal derecha y abocamiento ectópico del uréter en dilatación quística de la vesícula seminal. *Actas Urológicas Españolas* 2004;28:688-93.
79. Madrid García FJ, Madroño Cuevas C, Rivas Escudero JA, et al. Seminal vesicle cyst and ipsilateral renal agenesis. *Arch Esp Urol* 2004;57:168-71.
80. Kosan M, Tul M, Inal G, et al. A large seminal vesicle cyst with contralateral renal agenesis. *Int Urol Nephrol* 2006;38:591-2.
81. Gözen AS, Alagöl B. Endoscopic management of seminal-vesical cyst with right renal agenesis causing acute urinary retention: case report. *J Endourol* 2006;20:919-22.
82. Romero Selas E, Montes M, Barbagelata A, et al. Seminal vesicle cyst and ipsilateral renal agenesis. *Arch Esp Urol* 2006;59:750.
83. Pal D, Bag A, Sarkar S. Giant seminal vesicle cyst with ipsilateral hypoplastic kidney: Report of a case with review of literature. *Indian J Urol* 2006;22:64-5.
84. Cihan A, Cimen S, Secil M, et al. Congenital seminal vesicle cyst accompanying ipsilateral renal agenesis and rudimentary ureter. *Int Urol Nephrol* 2006;38:133-5.
85. Chen HW, Huang SC, Li YW, et al. Magnetic resonance imaging of seminal vesicle cyst associated with ipsilateral urinary anomalies. *J Formos Med Assoc* 2006;105:125-31.
86. Suárez Aliago B, Rodríguez Jiménez I. Seminal vesicle cyst with renal agenesis associated with vascular anomalies. Case report. *Arch Esp Urol* 2007;60:1.134-6.
87. Han P, Dong Q, Shi M, et al. Seminal vesicle cyst and ipsilateral renal agenesis: Laparoscopic approach. *Arch Androl* 2007;53:285-8.
88. Carbone A, Palleschi G, Tomiselli G, et al. Renal aplastic dysplasia and ipsilateral ectopic ureter obstructing the seminal via: a possible cause of male infertility. *Eur Urol* 2007;52:268-72.
89. Labanaris AP, Zugor V, Meyer B, et al. A case of a large seminal vesicle cyst associated with ipsilateral renal agenesis. *ScientificWorldJournal* 2008;8:400-4.
90. Casey RG, Stunell H, Buckley O, et al. A unique radiological pentad of mesonephric duct abnormalities in a young man presenting with testicular swelling. *Br J Radiol* 2008;81:e93-6.
91. Stimac G, Dimanovski J, Spajic B, et al. Seminal vesicle cyst with ectopic ureteral insertion and ipsilateral renal dysplasia--an unexpected diagnosis. *Acta Clin Croat* 2008;47:101-4.
92. Ohgaki K, Horiuchi K, Oka F, et al. A case of seminal vesicle cyst associated with ipsilateral renal agenesis diagnosed during an investigation of perineal pain. *J Nippon Med Sch* 2008;75:122-6.
93. Selli C, Cavalleri S, De Maria M, et al. Robot-assisted removal of a large seminal vesicle cyst with ipsilateral renal agenesis associated with an ectopic ureter and a müllerian cyst of the vas deferens. *Urology* 2008;71:1226.e5-7.
94. Sotelo R, Herrera C, Carmona O, et al. Robot-assisted excision of a Mullerian cyst with anastomosis of the vas deferens to the seminal vesicle. *J Urol* 2009;74:S163-4.
95. Pereira BJ, Sousa L, Azinhais P, et al. Zinner's syndrome: an up-to-date review of the literature based on a clinical case. *Andrologia* 2009;41:322-30.
96. Seo IY, Kim HS, Rim JS. Congenital seminal vesicle cyst associated with ipsilateral renal agenesis. *Yonsei Med J* 2009;50:560-3.
97. Ghonge NP, Aggarwal B, Sahu AK. Zinner syndrome: a unique triad of mesonephric duct abnormalities as an unusual cause of urinary symptoms in late adolescence. *Indian J Urol* 2010;26:444-7.
98. Lim CS, Lee DE, Mackie S, et al. Seminal vesicle cyst, vas deferens ectopia, pelvic kidney and cryptorchidism. *ANZ J Surg* 2010;80:382.
99. Viktorsdóttir MB, Jónsson E, Einarisdóttir H. Case report: Seminal vesicle cyst with ipsilateral renal agenesis. *Laeknabladid* 2010;96:619-20.
100. Allaparthi S, Blute RD Jr. Novel application of da Vinci robotic system in patients of Zinners syndrome--case report and review of literature. *Can J Urol* 2010;17:5109-13.
101. Dombóvári P, Mavrogenis S, Borka K, et al. Seminal vesicle cyst presenting as pelvic abscess. *Journal of Sexual Medicine* 2010;7:458.
102. Ahallal Y, Tazi MF, Khallouk A, et al. Conservative management of a congenital seminal vesicle cyst associated with ipsilateral renal agenesis revealed by cystitis: one case report. *Case Rep Urol* 2011;2011:125753.
103. Hampton LJ, Moore BW, Orton VA, et al. Robotic excision of a large seminal vesicle cyst: Zinners syndrome. *J Endourol* 2011;25:A326.
104. Kovac JR, Golev D, Khan V, et al. Case of the month # 168: seminal vesicle cysts with ipsilateral renal dysgenesis. *Can Assoc Radiol J* 2011;62:223-5.
105. Jang KD, Choi KH, Yang SC, et al. Laparoendoscopic single-site surgery (LESS) for excision of a seminal vesicle cyst associated with ipsilateral renal agenesis. *Korean J Urol* 2011;52:431-3.
106. Kalyanaraman B, Shah AK, Vemana G, et al. Seminal vesicle cyst as an unusual cause of anejaculation. *J Endourol* 2012;26:A223.
107. Devaraju S, Nerli RB, Hiremath MB. A large mullerian duct cyst presenting as an abdominal mass with ipsilateral renal agenesis: an unusual presentation. *Nephrourol Mon* 2012;4:640-3.
108. Basal S, Alp BF, Demirel Z, et al. Robot assisted laparoscopic surgery for seminal vesicle cyst. *J Endourol* 2012;26:A220-1.
109. El-Assmy A, Abou-El-Ghar ME. Giant seminal vesicle cyst causing ipsilateral hydronephrosis with contralateral renal agenesis. *Urology* 2012;79:e17-8.
110. Haghighi R, Shadpour P, Maghsudy R. Renal agenesis associated with ipsilateral ectopic ureter entering a large seminal vesicle. *J Endourol* 2012;26:A431.
111. Coşkun B, Dalkılıç A, Sönmez NC, et al. A case of seminal vesicle cyst associated with ipsilateral renal agenesis diagnosed during an investigation of urinary incontinence. *Türk J Urol* 2013;39:53-5.
112. Kardoust Parizi M, Shakhssalim N. Management of Zinner's syndrome associated with contralateral seminal vesicle hypoplasia: a case report. *Case Rep Urol* 2013;2013:494215.
113. Altobelli E, Bove AM, Falavolti C, et al. Robotic-assisted laparoscopic approach in the treatment for Zinner's Syndrome associated with ipsilateral megaureter and incomplete double-crossed ectopic ureter. *Int Urol Nephrol* 2013;45:635-8.
114. Silveri M, Zaccara A, Colajacomo M, et al. ... And if this was a Zinner's syndrome? *J Pediatr Surg Case Rep* 2014;2:530-2.
115. Batur A, Karakose S, Karalezli G. Congenital seminal vesicle cyst accompanying ipsilateral renal and ureteral agenesis. *Urol J* 2014;11:1203.
116. Palmer WC, Patel NC, Renew JR, et al. Acute infection of a documented seminal vesicle cyst via hematogenous seeding.

- Urol J 2014;10:1157-9.
117. Lee KH, Liao ACH, Liu CL, et al. Laparoscopic excision of a huge seminal vesicle cyst associated with ipsilateral renal agenesis. *J Endourol* 2014;28:A306.
 118. Shepherd G, Rajimwale A. Embryology of the absent vas supported by 2 cases of congenital unilateral absence of vas with varied associations. *Urol Case Rep* 2014;2:49-50.
 119. Kanavaki A, Vidal I, Merlini L, et al. Congenital seminal vesicle cyst and ipsilateral renal agenesis (Zinner syndrome): a rare association and its evolution from early childhood to adolescence. *European J Pediatr Surg Rep* 2015;3:98-102.
 120. Burak Özkan M, Ceyhan Bilgici M, Şahin M, et al. Congenital seminal vesicle cyst accompanying with ipsilateral renal agenesis in an adolescent patient: a pediatric radiologist approach to Zinner's syndrome. *The Egyptian Journal of Radiology and Nuclear Medicine* 2015;46:1159-62.
 121. Donato P, Roberts MJ, Hirst J, et al. Acute obstructive nephropathy in a solitary kidney secondary to massive seminal vesicle cyst. *ANZ J Surg* 2018;88:E345-7.
 122. Alp B, Demirer Z, Guragac A, et al. Ipsilateral Renal Agenezinin Eslik Ettigi Dev Seminal Vezikul Kistinin Robot Yardimli Laparoskopik Cerrahi Ile Tedavisi. *Gülhane Tıp Derg* 2015;57:331-4.
 123. Kim Y, Baek HW, Choi E, et al. Squamous cell carcinoma of the seminal vesicle from zinner syndrome: a case report and review of literature. *J Pathol Transl Med* 2015;49:85-8.
 124. Pavan N, Bucci S, Mazzon G, et al. It's not always varicocele: a strange case of Zinner syndrome. *Can Urol Assoc J* 2015;9:E535-8.
 125. Juho YC, Wu ST, Tang SH, et al. An unexpected clinical feature of Zinner's syndrome - a case report. *Urol Case Rep* 2015;3:149-51.
 126. García Asensio D, Fernández Martín M. Zinner syndrome diagnosed in a child. *Radiologia* 2016;58:73-4.
 127. Aghaways I, Ahmed SM. Endourologic intervention for management of infertility in a man with Zinner syndrome resulting in a natural pregnancy. *J Endourol Case Rep* 2016;2:71-3.
 128. Hajji F, Hammoune N, Miloudi M, et al. Unusual variant of inverted Y ureteral duplication with an ipsilateral seminal vesicle cyst and renal dysgenesis. *Ann R Coll Surg Engl* 2017;99:e19-21.
 129. Dogan F, Xhafa A, Bimbashi B. Excision of a giant seminal vesicle cyst associated with ipsilateral renal agenesis Zinner's Sendrom. *J Endourol* 2016;30:A417.
 130. Shah S, Patel R, Sinha R, et al. Zinner syndrome: an unusual cause of bladder outflow obstruction. *BJR Case Rep* 2017;3:20160094.
 131. Bakloul F, Jakhlal N, Elghazoui A, et al. Seminal vesicle cyst: about a case. *Pan Afr Med J* 2016;25:68.
 132. Gutierrez García MA, Estebanez Zarranz MJF, Crespo Crespo I, et al. Zinner syndrome, robotic management. *European Urology Supplements* 2016.
 133. Maré A, Liddell H, Chan HF. Ectopic ureteric insertion into a seminal vesicle cyst. A case report of recurrent UTI's and an unusual fistulous tract. *BJU Int* 2015;115:76.
 134. Slaoui A, Regragui S, Lasri A, et al. Zinner's syndrome: report of two cases and review of the literature. *Basic Clin Androl* 2016;26:10.
 135. Canales-Casco N, Dominguez-Amillo A, Arrabal-Polo MA, et al. Hematospermia as a rare form of presentation of Zinner syndrome. *Urology* 2017;99:e15-6.
 136. Prem G, Goh EH, Fam XI, et al. A rare presentation of urinary tract symptoms in a teenager: Zinner's syndrome. *BJU Int* 2017;120:33.
 137. Chiapparrone G, Liguori G, Pavan N, et al. Fertility recovery after laparoscopic removal of hypertrophic seminal vesicle. *J Urol* 2017;197:e1060.
 138. Cao JD, Zhu SL, Bai ZG, et al. Zinner syndrome: a case report and review of the literature. *Zhonghua Nan Ke Xue* 2017;23:452-4.
 139. Ahmed A. Ipsilateral renal agenesis with megaureter, blind end proximal ureter and ureterocele in an adult. *J Ayub Med Coll Abbottabad* 2017;29:150-3.
 140. Campora M, Olivero A, Toncini C, et al. Zinner syndrome: a diagnostic challenge. The aid of morphology, embryology, and immunohistochemistry. *Urology* 2017;108:e3-5.
 141. Khanduri S, Katyal G, Sharma H, et al. Unique association of multiple seminal vesicle cysts with contralateral renal agenesis: a rare variant of Zinner syndrome. *Cureus* 2017;9:e1415.
 142. Roth C, Götz G, Gräfe D, et al. Zinner syndrome: underestimated differential diagnosis in children with groin pain and retrovesical cyst. *Pediatric Radiology* 2018;48:S581-2.
 143. Millman AL, Krocak T, Ordon M. Images - Large symptomatic seminal vesicle cyst treated with robotic-assisted seminal vesiculectomy. *Can Urol Assoc J* 2019;13:E99-101.
 144. Razdan S, Kryvenko ON, Razdan S. Robotic-assisted laparoscopic vesiculectomy in a patient with atypical Zinner syndrome presenting with large cyst involving bilateral seminal vesicles and vasa deferentia. *Urol Case Rep* 2018;18:79-81.
 145. Florim S, Oliveira V, Rocha D. Zinner syndrome presenting with intermittent scrotal pain in a young man. *Radiol Case Rep* 2018;13:1224-7.
 146. Kord E, Zisman A, Darawsha AE, et al. Minimally invasive approach for treatment of seminal vesicle cyst associated with ipsilateral renal agenesis. *Urol Int* 2017;99:338-42.
 147. Farooqui A, AlDhahir L, Mahfooz AB. Massive seminal vesicle cyst with ipsilateral renal agenesis - Zinner syndrome in a Saudi patient. *Urol Ann* 2018;10:333-5.
 148. Brown M, Aiken W. Case report: Zinner's syndrome and a blind ending obstructed ureterocele: an unusual cause of lower urinary tract symptoms in a young man. *BJU Int* 2018;121:11-2.
 149. El Mortaji H, Elatiqi K, El Hammaoui H, et al. Zinner's syndrome: a case report. *Prog Urol* 2018;28:464-5.
 150. Briosia F, Valsassina R, Mira C, et al. Zinner and Mayer-Rokitansky-Küster-Hauser syndromes: when unilateral renal agenesis meets genital anomalies. *BMJ Case Rep* 2019;12:e229034.
 151. Symeonidis EN, Gkekas C, Tsifountoudis I, et al. Incidental finding of Zinner syndrome in a Greek military recruit: a case report of a rare clinical entity. *Mil Med Res* 2019;6:4.
 152. Kori R, Bains L, Lal P, et al. Zinner syndrome mimicking bladder outlet obstruction managed with aspiration. *Urol Ann* 2019;11:449-52.
 153. AlArifi M, Al-Gahwary M, Gomha M. The association of renal agenesis and ipsilateral seminal vesicle cyst: Zinner syndrome case report. *Case Rep Urol* 2019;2019:1242149.
 154. Aslan S. A rare cause of chronic pelvic pain in young man: magnetic resonance imaging findings of Zinner's syndrome. *Journal of Urological Surgery* 2019;6:331-4.
 155. Cito G, Sforza S, Gemma L, et al. Infertility case presentation in Zinner syndrome: can a long-lasting seminal tract obstruction cause secretory testicular injury? *Andrologia* 2019;51:e13436.
 156. Bhukte S. Robot assisted laparoscopic excision of Seminal vesicle cyst: a rare entity (Zinner syndrome: Unilateral renal agenesis, ipsilateral seminal vesicle cyst and ejaculatory duct obstruction). *BJU Int* 2019;123:47-8.
 157. Patil M, Shah R, Shah S, et al. Unusual variant of zinner syndrome presenting as cause of infertility in young male. *Indian J Urol* 2019;35:S31-2.
 158. Sahare P, Shah S, Singh T, et al. A rare case of Zinn's syndrome presenting as LUTS in young adult managed laparoscopically. *Indian J Urol* 2019;35:S31.
 159. AlhajerI F, Salem S, Harraz A, et al. Recurrent epididymo-orchitis: uncommon presentation of zinner syndrome. *J Endourol* 2019;33:A370.
 160. Corongiu E, Grande P, Olivieri V, et al. Minimally invasive management of a symptomatic case of Zinner's syndrome: Laparoscopic seminal vesiculectomy and ipsilateral nephroureterectomy. *Arch Ital Urol Androl* 2019;91:58-9.
 161. Chibeleian BC, Tătaru OS, Borda A, et al. Zinner's syndrome – the value of clinical imaging and morphopathological findings for diagnosis. *Journal of Interdisciplinary Medicine* 2019;4:196-200.
 162. Bryson CF, Delpe S, Tatzel S, et al. Robot-assisted excision of congenital mega-seminal vesicle associated with Zinner's syndrome. *J Endourol Case Rep* 2019;5:4-6.
 163. Hergan B, Fellner FA, Akbari K. Incidental imaging findings suggesting Zinner syndrome in a young patient with pulmonary embolism: a case report. *Radiol Case Rep* 2020;15:437-41.
 164. Yadav SP, Gurjar K. Zinner syndrome: an unusual cause of bladder outflow obstruction. *Indian J Urol* 2020;36:S35.
 165. Tan Z, Li B, Zhang L, et al. Classifying seminal vesicle cysts in the diagnosis and treatment of Zinner syndrome: a report of six cases and review of available literature. *Andrologia* 2020;52:e13397.
 166. Jatal S. Say hi to zinner - Laparoscopic seminal vesical cyst excision. *Indian J Urol* 2018;34:40.